

(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 773 020 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication
14.05.1997 Bulletin 1997/20

(51) Int Cl⁶: **A61K 31/205, A61K 31/22,
A61K 31/05**

(21) Application number: **96830517.7**

(22) Date of filing: **11.10.1996**

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**

(30) Priority 17.10.1995 IT RM950687

(71) Applicant: **Sigma-Tau Industrie
Farmaceutiche Riunite S.p.A.
I-00144 Roma (IT)**

(72) Inventor: **Cavazza, Claudio
00186 Roma RM (IT)**

(74) Representative: **Cavattoni, Fabio et al
Cavattoni & Raimondi
Viale dei Parioli, 160
00197 Roma (IT)**

(54) **Pharmaceutical compositions containing L-carnitine or derivatives thereof in combination with resveratrol or derivatives thereof, for the prophylaxis and treatment of cardiovascular disorders, peripheral vascular diseases and peripheral diabetic neuropathy**

(57) The co-ordinated therapeutic use of L-carnitine, lower alkanoyl L-carnitines or the pharmacologically acceptable salts thereof with resveratrol, resveratrol derivatives or resveratrol-containing natural products is

disclosed for producing a medicament for the prophylaxis and treatment of cardiovascular disorders, peripheral vascular diseases and peripheral diabetic neuropathy.

EP 0 773 020 A2

Description

The present invention relates to a novel therapeutic use of L-carnitine, lower alkanoyl L-carnitine and their pharmacologically acceptable salts

More specifically, the present invention relates to the co-ordinated use of L-carnitine, alkanoyl L-carnitines or their pharmacologically acceptable salts and a trihydroxy or tetrahydroxystilbene for the prophylaxis and treatment of cardiovascular diseases, peripheral vascular diseases and diabetic peripheral neuropathy.

Within the scope of the present invention, by "co-ordinated use" of the aforesaid compounds it is meant indifferently both the coadministration, i.e. the substantially concomitant supplementation of L-carnitine or alkanoyl L-carnitine or a pharmacologically acceptable salt thereof and a trihydroxy or tetrahydroxystilbene as active ingredients, and the administration of a combination preparation containing a mixture of the aforesaid active ingredients, in addition to suitable excipients, if any.

Consequently, the present invention also relates to orally, parenterally, rectally or transdermally administrable pharmaceutical compositions suitable for the treatment of cardiovascular disorders, peripheral vascular diseases and diabetic peripheral neuropathy, which comprises L-carnitine or a lower alkanoyl L-carnitine or a pharmaceutically acceptable salt thereof and a trihydroxy or tetrahydroxystilbene as active ingredients.

Among the trihydroxy and tetrahydroxystilbenes, resveratrol, 3,4',5-trihydroxystilbene, the derivatives thereof and the resveratrol-containing natural products, are particularly preferred.

Resveratrol and derivatives thereof are available as extracts or powders of natural products, mainly extracted from vitis vinifera and particularly from the skin, grapes, grape-seeds, grape-stalks and leaves of grapevine.

The alkanoyl L-carnitines useful for the novel therapeutic use of the present invention are those wherein the alkanoyl group, straight or branched, has 2 to 8, preferably 2 to 6, carbon atoms.

Particularly preferred are acetyl, propionyl, butyryl, valeryl and isovaleryl L-carnitine.

Pharmaceutically acceptable salts of L-carnitine or alkanoyl L-carnitine include, in addition to the inner salts, all pharmaceutically acceptable salts which are prepared by the addition of an acid to L-carnitine or alkanoyl L-carnitine, respectively, and which do not give rise to undesirable toxic or collateral effects. The formation of pharmaceutically acceptable acid addition salts is well known in the pharmaceutical technology.

Non-limiting examples of suitable salts include the chloride, bromide, orotate, acid aspartate, acid citrate, acid phosphate, acid fumarate, lactate, malate and acid malate, acid oxalate, acid sulphate, glucose phosphate, tartrate and acid tartrate.

For the sake of simplicity and clarity, hereinbelow reference will be made to L-carnitine and resveratrol and

its derivatives only, it being understood, however, that whatever disclosed in connection with L-carnitine and resveratrol equally applies to the above-identified alkanoyl L-carnitines and the pharmacologically acceptable salts of both L-carnitine and the foregoing alkanoyl L-carnitines as well as to trihydroxy and tetrahydroxystilbenes, in general.

Previous therapeutic uses of L-carnitine are already known. For instance, L-carnitine has been used in the cardiovascular field in the treatment of acute and chronic myocardial ischaemia, angina pectoris, cardiac arrhythmias and insufficiency. In nephrology, L-carnitine has been administered to chronic uraemic patients who are subject to regular haemodialysis treatment with a view to counteracting muscular asthenia and the onset of muscular cramps. Further therapeutic uses relate to the restoration to normal of unbalanced HDL/(LDL+VLDL) ratio and total parenteral nutrition. It is also known the use of L-carnitine for treating certain myopathies and muscular dystrophies.

Resveratrol, 3,4',5-trihydroxystilbene, belongs to the class of stilbenic phytoalexins which include various trihydroxy- and tetrahydroxystilbenes.

Resveratrol is the one present in largest amounts in red grapes and responsible for favourable pharmacological effects.

Data have recently been reported on the biological activity of resveratrol (Elliot G.R., Lanwen A.P.M., Bonta L.L., Agents Actions 32, 88, 1991; Ronca F., Palmieri L., Malengo S., Bertelli A., Int. J. Tiss. Reac. 16, 187, 1994; Amico Roxas M., Caruso A., Cutuli V.M.C., Bernardis E., INWIN ABS-Geneva 31-4-1993).

In popular medicine there was already an awareness that extracts from the roots of plants (*Polygonum cuspidatum* L. and *Polygonum multiflorum*) containing products such as resveratrol could be used in the prevention and therapy of atherosclerosis (Arichi H., Kimura Y., Okuda H., Baba K., Kozawa M., Arichi S., Chem. Pharm. Bull., 30, 1766, 1982).

More recently, this activity of theirs has been found to be related mainly to resveratrol and its ability to inhibit cyclo-oxygenases and lipo-oxygenases with a consequent reduction in the production of thromboxanes and leukotrienes which are known for their inflammatory and platelet aggregation action (Kimura Y., Okuda H., Arichi S., Biochem. Biophys. Acta 834, 275, 1985; Kimura Y., Okuda H., Arichi S., J. Med. Pharm. Soc. Watran-Yaku 2, 516, 1985) and to its antihyperoxidative ability (Kimura Y., Ohminami H., Okuda H., Baba K., Kozawa M., Arichi S., Plant. Med. J. Med. Plant Res. 49, 51, 1983).

Recent data in the literature indicate, moreover, that the onset of diabetic peripheral neuropathy is facilitated by increased platelet aggregation [(Medical treatment of diabetic arteriopathy) Got I., Ziegler J., Drouin P., J. Med. Vasc. 18(1), 30-6, 1983. (Coagulation and fibrinolytic parameters in patients with various angiopathies - analysis in cerebral thrombosis, diabetic and vasculitic neuropathies) Kohriyama T., Katayama S., Tanaka E.,

Yamamura Y., Nakamura S., *Rinsho-Shinkeigaku Jun* 33(6), 696-11, 1993; (Relationships between haemostatic factors and capillary morphology in human diabetic neuropathy) Ford I., Malik R.A., Newrick P.G., Preston F.E., Ward J.D., Greaves M., *Thromb. Haemost.* Dec 7 68(6), 628-33, 1992].

There has recently been an increase in interest in resveratrol as a result of epidemiological data showing a lower incidence of mortality due to cardiovascular damage even in populations with a high calorie consumption with a high percentage of lipids, but whose diets included red wine, as compared to populations who had a lower calorie consumption and lower percentage of lipids, but whose diets did not include wine, particularly red wine (Seigneur M., Bonnet Y., Dorian B., Beuchimal D., Dronillet F., Gouverneur G., Larrue Y., Crockett R., Bricaud H., *J. Appl. Card.* 5, 215, 1990; Siemann E.H., Creasy L.L., *Am. J. Enol. Vitic.* 43, 49, 1992; Renaud S., De Lorgeril M., *Lancet* 339, 1523, 1992; Scharp D., *Lancet* 341, 27, 1993).

Research conducted on the components present in red wine possibly responsible for this favourable effect has succeeded in identifying the presence in the wine of numerous products from grapes which are endowed with pharmacological properties, and particularly the presence of high percentages of resveratrol (Creasy L.L., Coffee L.L., *J. Am. Soc. Hort. Sci.* 113, 230, 1988; Ylandet P., Bessis R., Ganoixheron B., *Am. J. Enol. Vitic.* 42, 41, 1991).

Its isolation and then its synthesis have revealed that this phytoalexin effectively possesses many pharmacological activities which can be regarded as responsible for the protective effects of red wine at cardiovascular level (Frankel E.N., Kanner Y., Parks E., Kinsella Y.E., *Lancet* 341, 454, 1993).

In addition to its ability to modulate the synthesis of eicosanoids, resveratrol has proved capable of promoting the formation of nitroxides with a vasodilatory action and of inhibiting platelet aggregation induced by collagen or ADP (Fitzpatrick D., Steven L., Coffee R.G., *Am. J. Physiol.*, 265 (Heart Circ. Physiol. 34, 774, 1993).

The resveratrol content of red wine is related to the type of grape the wine is produced from and the region where the grapes are grown (Mattiviti F., *Riv. Vitic. Enol.* 1, 37, 1993).

Its presence, in differing percentages, can also be found both in the grape skins and in the pips or grape stalks (Langcake P., Pryce R., *Experimentia* 33, 151, 1977; Pezet R., Pont V., *Plant Physiol. Biochem.* 25, 603, 1988).

Its concentration is higher in grapes affected by a number of typical diseases of the vine and this has been attributed, amongst other things, to the antimould and antiparasite effect which resveratrol is capable of exerting and to an increase in its productive self-regulation in the presence of such diseases (Dercks W., Creasy L.L., *Physiol. Molec. Plant Pathol.* 34, 189, 1989; Langcake P., *Physiol. Plant Pathol.* 18, 213, 1981).

The invention described herein is based on the surprising synergic effect which occurs on combining carnitine or one of its derivatives and resveratrol or one of its derivatives, or natural compounds containing resveratrol.

The compositions according to the invention prove particularly effective in inhibiting platelet aggregation, vascular thrombo-embolic lesions and atherosclerosis and can thus be used in the pharmaceutical, dietetic and alimentary fields for the prevention or treatment of cardiovascular diseases, peripheral cardiopathies and diabetic peripheral neuropathies.

In such compositions the weight-to-weight ratio of resveratrol or its derivatives or extracts or powder of grapes containing resveratrol to L-carnitine or its derivatives ranges from 1:1 to 1:1000 and is preferably 1:500.

On the one hand, in fact, L-carnitine and particularly propionyl carnitine can act by varying the lipid substrate from which come, as a result of the action of cyclo-oxygenases and lipo-oxygenases, the various vasoconstricting and pro-aggregation factors, reducing their formation and facilitating the synthesis of antiaggregating and vasodilatory factors.

Resveratrol, on the other hand, can directly inhibit the action of the cyclo-oxygenase and lipo-oxygenase enzymes and the release of pro-inflammatory eicosanoids and leukotrienes and platelet aggregation promoters. To this we should add the antihyperoxidative effect common both to L-carnitine and to resveratrol and the ability of carnitine to inhibit the vasoconstricting and hypertensive action of endothelin, whereas resveratrol may be capable of releasing nitroxide which is known to exert a vasodilating action.

The routes of interaction between carnitine and resveratrol are therefore multiple and a combination of the two substances is useful for the purposes of preventing and treating pathological conditions related to cardiovascular disorders of the thrombo-embolic and atherosclerotic type and to diabetic peripheral neuropathy. We shall now report the results of a number of tests which confirm the unexpected potent synergism between carnitine and resveratrol or extracts or powders from the skins, pips or stalks of grapes and containing high percentages of resveratrol.

Toxicology

The tests performed both in mice and in rats administering combinations of high doses both of L-carnitine or its derivatives and resveratrol or grape extracts (with a resveratrol concentration of approximately 0.1%) have demonstrated the low toxicity and good tolerability of the new composition. The administration of up to 1 g/kg of L-carnitine or its derivatives per os in the rat together with high doses of resveratrol (up to more than 0.5 g/kg) or grape extracts (up to more than 1 g/kg) revealed no toxic reactions worthy of note. Similarly well tolerated appeared to be protracted administration through the di-

et every day for three consecutive months with a combination of both L-carnitine and its derivatives together with resveratrol or grape extracts containing resveratrol. At the end of this treatment no abnormalities of any of the various biological parameters considered (weight gain, haematocrit, serum glucose, BUN, etc.) were detected, nor of the main organs examined at histology (liver, heart, lungs, adrenal glands, gonads)

Platelet aggregation tests

Platelet-rich plasma (PRP) was obtained from the blood of volunteers by centrifuging the plasma at 145 x g for 5 min at 22°C. The number of platelets was determined using a CA 580A (Delcon) Platelet Counter. The number of platelets was brought down to a fixed value of 300.000 platelets/ml by adding platelet-poor plasma (PPP). PPP was obtained by centrifuging blood at 1600 x g for 10 min at 22°C.

Platelet aggregation was determined photometrically (Born G.V.R., Nature 194, 927, 1962) using an Elvi 840 aggregometer in plastic test tubes under continual stirring at 1000 rpm. The maximum light transmission (100%) was measured with PPP readings, whereas 0% transmission was measured with PRP readings.

The degree of aggregation of PRP induced by the aggregating agent (collagen at 2.5 to 5 ng/ml) was calculated as a percentage of maximum transmission of light obtained with PPP using 250 1 PRP samples.

Aggregation was measured in basal conditions and after 10 minutes of incubation with L-carnitine, resveratrol, grape extract, and combinations of these preparations. Inhibition of the platelet aggregation induced by collagen (2.5 ng/ml) proved evident (ED₅₀ 3.5 ng/ml) for resveratrol and for grape extract (ED₅₀ with a resveratrol concentration equal to 2.5 ng/l), whereas for carnitine or its derivatives there was no significant change. However, when using a combination of the carnitines plus resveratrol at the same doses, 100% inhibition of platelet aggregation was achieved, thus showing a marked synergism between L-carnitine and resveratrol or grape extract containing resveratrol.

Tests with experimental atherosclerosis

A potentiation of the joint action of L-carnitine and resveratrol or grape extract has also been demonstrated in tests involving the induction of experimental atherosclerosis. The atherosclerotic vascular lesions induced according to the method described by M.R. Malinow (Atherosclerosis 48, 105, 1983) by administering an atherogenic diet (24% casein, 10% cotton oil, 5% salt, 60% sugar, 1% cholesterol, vitamin D₂ 200 m STU/g diet) to different groups of male Wistar rats for 6 consecutive weeks were reduced both by carnitine and by resveratrol, but the reductions were surprising when these products were administered in combination. With the administration of propionyl L-carnitine and resveratrol

atherosclerotic lesions at the level of the aorta were completely inhibited, thus demonstrating an evident synergic effect of carnitine and resveratrol.

By way of examples and without in any way limiting the possibilities to those listed we give here a number of the possible formulations related to the invention described herein

Examples

- 1) L-carnitine 500 mg, resveratrol 5 mg;
- 2) acetyl L-carnitine 500 mg, resveratrol 5 mg;
- 3) propionyl L-carnitine 500 mg, resveratrol 5 mg;
- 4) isovaleryl L-carnitine 500 mg, resveratrol 5 mg;
- 5) valeryl L-carnitine 500 mg, resveratrol 5 mg;
- 6) butyryl L-carnitine 500 mg, resveratrol 5 mg;
- 7) L-carnitine 500 mg, lyophilized dry grape extract 250 mg (0.5% mg resveratrol titre);
- 8) acetyl L-carnitine 500 mg, lyophilized dry grape extract 250 mg (0.5% mg resveratrol titre);
- 9) propionyl L-carnitine 500 mg, lyophilized dry grape extract 250 mg (0.5% mg resveratrol titre);
- 10) isovaleryl L-carnitine 500 mg, lyophilized dry grape extract 250 mg (0.5% mg resveratrol titre);
- 11) L-carnitine 500 mg, resveratrol 10 mg;
- 12) L-carnitine 500 mg, lyophilized grape extract 250 mg (0.5% mg resveratrol titre), tocopherol acetate 20 mg, beta-carotene 10 mg;
- 13) L-carnitine 500 mg, resveratrol 5 mg, tocopherol acetate 20 mg, vitamin C 100 mg, selenium 40 mg.

Claims

1. Use of L-carnitine and alkanoyl L-carnitine wherein the alkanoyl group, straight or branched, has 2-8 preferably 2-6, carbon atoms or the pharmacologically acceptable salts thereof and a trihydroxy or tetrahydroxystilbene for producing a pharmaceutical composition for the prophylaxis and treatment of cardiovascular disorders, peripheral vascular diseases and diabetic peripheral neuropathy
2. The use of claim 1, wherein the trihydroxystilbene is selected from the group consisting of resveratrol the glucoside thereof 3,4',5'-trihydroxystilbene-3-beta-mono-D-glucoside, and the pharmacologically acceptable salts and esters thereof.
3. The use of claim 1 or 2, wherein the alkanoyl L-carnitine is selected from acetyl, propionyl, butyryl, valeryl and isovaleryl L-carnitine
4. An orally, parenterally, rectally or transdermally administrable pharmaceutical composition for treating cardiovascular disorders, peripheral vascular diseases and diabetic peripheral neuropathy which comprises L-carnitine or an alkanoyl L-carnitine

wherein the alkanoyl group, straight or branched, has 2-8, preferably 2-6, carbon atoms or the pharmacologically acceptable salts thereof and a trihydroxy or tetrahydroxystilbene as active ingredients and a pharmacologically acceptable excipient thereof 5


5. The composition of claim 4 wherein the trihydroxy or tetrahydroxystilbene is selected from the group consisting of 3,4',5-trihydroxystilbene (resveratrol), the glucoside thereof, 3,4',5-tri-hydroxystilbene-3-beta-mono-D-glucoside, and the pharmacologically acceptable salts and esters thereof 10
6. The composition of claim 4 or 5, wherein the alkanoyl L-carnitine is selected from acetyl, propionyl, butyryl, valeryl and isovaleryl L-carnitine 15
7. The composition of claims 4-6 which comprises resveratrol-containing natural products, extracts and powders from *vitis vinifera*. 20
8. The composition of claim 7 wherein the weight ratio between resveratrol, derivatives thereof or resveratrol-containing extracts and L-carnitine or said alkanoyl L-carnitine is from 1:1 to 1:1000, preferably 1:500. 25
9. The composition of claims 4-8 which further comprises polyphenols, anthocyanins, anthocyanosides, vitamins, mineral salts, antioxidant agents and vegetal fibers. 30
10. The composition of claims 4-9 as a solid, semisolid, liquid, semiliquid composition in the form of powders, granules, liposomes, tablets, capsules and vials 35

40

45

50

55



(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 773 020 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3
10.12.1997 Bulletin 1997/50

(51) Int Cl⁶ **A61K 31/205, A61K 31/22,
A61K 31/05**

(43) Date of publication A2
14.05.1997 Bulletin 1997/20

(21) Application number **96830517.7**

(22) Date of filing **11.10.1996**

(84) Designated Contracting States:
**AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC
NL PT SE**

(30) Priority **17.10.1995 IT RM950687**

(71) Applicant: **Sigma-Tau Industrie
Farmaceutiche Riunite S.p.A.
00144 Roma (IT)**

(72) Inventor: **Cavazza, Claudio
00186 Roma RM (IT)**

(74) Representative: **Cavattoni, Fabio et al
Cavattoni & Raimondi
Viale dei Parioli, 160
00197 Roma (IT)**

(54) **Pharmaceutical compositions containing L-carnitine or derivatives thereof in combination with resveratrol or derivatives thereof, for the prophylaxis and treatment of cardiovascular disorders, peripheral vascular diseases and peripheral diabetic neuropathy**

(57) The co-ordinated therapeutic use of L-carnitine, lower alkanoyl L-carnitines or the pharmacologically acceptable salts thereof with resveratrol, resveratrol derivatives or resveratrol-containing natural products is

disclosed for producing a medicament for the prophylaxis and treatment of cardiovascular disorders, peripheral vascular diseases and peripheral diabetic neuropathy

EP 0 773 020 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 96 83 0517
shall be considered, for the purposes of subsequent
proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IntCl.6)
Y	EP 0 167 115 A (MAGIS FARMACEUTICI) 8 January 1986 * the whole document *	1-10	A61K31/205 A61K31/22 A61K31/05
Y	EP 0 256 999 A (SIGMA-TAU INDUSTRIE) 24 February 1988 * the whole document *	1-10	
Y	GB 2 043 443 A (CAVAZZA) 8 October 1980 * the whole document *	1-10	
Y	WANG Y-Z ET AL: "Reducing effect of 3,4',5-trihydroxystilbene-3-beta-mono-D-glucoside on arterial thrombosis induced by vascular endothelial injury" ACTA PHARMACOLOGICA SINICA, 16 (2). 1995. 159-162., XP002042342 * the whole document *	1-10	
-/-			
			TECHNICAL FIELDS SEARCHED (Int Cl.6)
			A61K
INCOMPLETE SEARCH The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims. Claims searched completely Claims searched incompletely Claims not searched Reason for the limitation of the search see sheet C			
Place of search		Date of publication of the search	Examiner
THE HAGUE		3 October 1997	Mair, J
CATEGORY OF CITED DOCUMENTS X particularly relevant if taken alone Y particularly relevant if combined with another document of the same category A technological background O non written disclosure P intermediate document		1 theory or principle underlying the invention 2 earlier patent document, but published on, or after the filing date 3 document cited in the application 4 document cited for other reasons 5 member of the same patent family, corresponding document	



European Patent
Office

INCOMPLETE SEARCH
SHEET C

Application Number
EP 96 83 0517

Claim(s) searched completely:
2,5

Claim(s) searched incompletely:
1,3,4,6-10

Reason for the limitation of the search:

The terms trihydroxy or tetrahydroxystilbene are not clear and concise definitions of chemical compounds. The search has had to be restricted to the specifically mentioned and claimed compounds and the general concept of the application.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 96 83 0517

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y	ZHANG P.-W. ET AL: "Influence of 3,4',5-trihydroxystilbene-3-beta-mono-D-glucoside on vascular endothelial epoprostenol and platelet aggregation" ACTA PHARMACOLOGICA SINICA, 1995, 16/3 (265-268), CHINA, XP002042343 * the whole document *	1-10	
Y	INAMORI Y ET AL: "THE BIOLOGICAL ACTIVITIES OF 3 4-O ISOPROPYLIDENE-3' 4 5'-TETRAHYDROXYSTILBENE" CHEM PHARM BULL (TOKYO), 33 (7), 1985, 2904-2909., XP002042344 * the whole document *	1-10	
Y	GRECH J N ET AL: "Novel Ca-2+-ATPase inhibitors from the dried root tubers of Polygonum multiflorum" JOURNAL OF NATURAL PRODUCTS (LLOYDIA), 57 (12), 1994, 1682-1687., XP002042345 * the whole document *	1-10	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
Y	DATABASE BIOSIS BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US LUO S-F ET AL: "INFLUENCES OF 3 4 5 TRIHYDROXYSTILBENE 3-BETA-MONO-D-GLUCOSIDE ON BEAT RATE AND INJURY OF CULTURED NEWBORN RAT MYOCARDIAL CELLS" XP002042349 * abstract * & ACTA PHARMACOL SIN, 11 (2), 1990, 147-150., --- -/--	1-10	

EPO FORM 1503 03 82 (P4C10)

PARTIAL EUROPEAN SEARCH REPORT

Application Number
EP 96 83 0517

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
Category	Citation of document with indication, where appropriate of relevant passages	Relevant to claim	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
Y	<p>PACE-ASCIK CR ET AL: "The red wine phenolics trans-resveratrol and quercetin block human platelet aggregation and eicosanoid synthesis: implications for protection against coronary heart disease."</p> <p>CLIN CHIM ACTA, MAR 31 1995, 235 (2) P207-19, NETHERLANDS, XP002042346</p> <p>* the whole document *</p> <p>---</p>	1-10	
Y	<p>GOLDBERG DM ET AL: "Beyond alcohol: beverage consumption and cardiovascular mortality."</p> <p>CLIN CHIM ACTA, JUN 15 1995, 237 (1-2) P155-87, NETHERLANDS, XP002042347</p> <p>* page 176, line 4-21 *</p> <p>---</p>	1-10	
D,A	<p>FORD, ISOBEL ET AL: "Relationships between haemostatic factors and capillary morphology in human diabetic neuropathy" THROMBOSIS AND HAEMOSTASIS, vol. 68, no. 6, 1992, pages 628-633, XP002042348</p> <p>* the whole document *</p> <p>-----</p>	1-10	